Psychological Services

Diagnostic Clinical Interview

| 90801 Diagnostic Interview Examination | |
|---|-------------|
| 90802 Interactive Diagnostic Interview Exam | |
| Student: | Date: |
| DOB: | School: |
| Age: | Grade: |
| Medicaid No.: | Student No. |
| Dx Code | |
| Source of Referral: | |
| Reason for Referral: | |
| Presenting Problem: | |
| Medical History: | |
| Current Medication: | |

| Previous Psychiatric/ Psychological Treatment / Intervention: | | |
|---|--|--|
| | | |
| Relevant Family History: | | |
| | | |
| | | |
| | | |
| Educational History: | | |
| | | |
| | | |
| | | |
| | | |
| History of alcohol or drug use: | | |

Mental Status Exam:

| Appearance: | □ Neat & well- | groomed | □Unk | empt | | | | |
|------------------------|-----------------|----------------|-------------------|-----------------|------------|-------------------|---------|---------|
| Responsiveness: | □ Alert | □ Distracted | | | | | | |
| Orientation: | □ X4 | □ Other | | | | | | |
| Attitude: | ☐ Cooperative | □ Passive | □G | uarded | ☐ Irrita | ble | | |
| Eye Contact: | ☐ Appropriate | □ Intense | □ P | oor | | | | |
| Motor Behavior: | □ Normal | ☐ Restless | □ Hy _] | peractive | | Hypoactive | | |
| Speech: | ☐ Spontaneous | □ Minim | nally res | ponsive | □ Slow | Į. | | |
| | □ Rapid | □ Slurred | | □Mumble | ed | | | |
| Vocabulary: | ☐ Average | | bove Av | erage | | Below Average | | |
| Thought Process: | □ Logical & go | al-directed | | □ Tar | ngential | | | |
| | □ Circumstantia | al | | □ Loc | ose Associ | ations | | |
| Thought Content: | □ Age-appropri | ate | ccupatio | ns 🗆 De | elusion | □ Compulsion | □ Phobi | as |
| Perception: | □ Normal | □ Auditory I | Hallucin | ations | □ Visua | al Hallucinations | 3 | |
| Suicidal: | □ Denied □ | Ideations | □ P | lan | □ Inten | tion | | |
| Homicidal: | □ Denied □ | Ideations | □ P | lan | □ Inten | tion | | |
| Mood: | □ Calm □ H | Нарру | □ Anxi | ous 🗆 | Depresse | d □ Irrita | able | □ Angry |
| Affect: | ☐ Appropriate | ☐ Inappropr | riate | □ Constri | cted | □ Labile | | |
| Insight: | □ Good | □ Fair | | □ Poor | | ☐ Externalizes | | |
| Judgement: □G | food | □ Fair | | Poor | | | | |
| Diagnostic Impression | on• | | | | | | | |
| 2 | | didi a a dh ad | . . | a f oods | of Clinica | .1 4444 | | |
| Axis I: Clinical Disor | raers/Otner coi | iaitions that | my be | a jocus o | of Cunica | u Attention- | | |
| Axis II: Personality I | Disorders/Men | tal Retardati | ion- (if | appropri | iate) | | | |
| Axis III: General Me | edical Conditio | ns- | | | | | | |
| Axis IV: Psychosocia | al and Environ | mental Prob | lems- | | | | | |
| Axis V: Global Asses | sment of Func | tioning- | Current | t | | Past Year | | |

Initial Clinical Evaluation Page 4 of 4

Suggested GAF ranges related to educational settings:

- 30 & below: hospitalized, residential treatment, specialized programs
- 30 40: violent, acting out
- 40 60: self-contained class, non-violent acting out, social problems
- 60 70: resource/itinerant services
- 70 & above: adjustment issues, probably not special education; guidance counseling

| Recommended Initi | ial Treatment / | Intervention | Plan: |
|-------------------|-----------------|--------------|-------|
|-------------------|-----------------|--------------|-------|

| - | Signature |
|---|-----------|
| | |